

QRS 1010 Pelvicenter

Repetitive peripheral magnetic stimulation to correct functional pelvic floor disorders

Scientific documentation and medical information

Women with Pelvic Pain Syndrome



Publisher:

© 2023 QRS International AG

Industriering 3 FL-9491 Ruggell Tel.: +423 798 80 88 / +31630916144 (WhatsApp)

E-Mail: emiel.spiessens@qrs-international.com / emiel.spiessens@gmail.com

YouTube: QRS GERMANY

QRS MedWiss Service
Reproduction prohibited

definition

Hardly any other clinical picture leads to an orthopedist as often and is misinterpreted as often as pain syndromes of the musculoskeletal system. The “pubic bone inflammation” stands out here in particular. But also the pelvic pain syndrome, postpartum pain, dyspareunia in women or the peritrochanteric pain that is often “diagnosed” as osteoarthritis of the hip seem to be somehow mysteriously connected. However, if one sees the pathomechanism of a myofascial syndrome behind it, important connections and thus therapy options open up.

Epidemiology and Prevalence

Osteitis pubis OB (pubic bone inflammation)

The frequency of OB in amateur and professional sports is between 0.5 and 7.0% [\[1\]](#), [\[2\]](#), [\[3\]](#), [\[4\]](#), [\[5\]](#). Although every sport is affected by OB (e.g. long-distance runners [\[6\]](#), [\[7\]](#)), it is much more common in sports with sprinting and kicking elements or quick changes of direction (soccer, American football, ice hockey, tennis and rugby) [\[8\]](#), [\[9\]](#), [\[10\]](#), [\[11\]](#). In addition to "spontaneous" healing processes, long-lasting symptoms that can last several months [\[12\]](#) and lead to long-term interruptions in training and competitions are not uncommon [\[13\]](#), [\[14\]](#).

Pelvic Pain Syndrome

According to data from a systematic review, the prevalence of female chronic pelvic pain syndrome (CPPS), which by definition lasts longer than 6 months, is between 5.7 and 26.6% [\[15\]](#). Female CPPS is different from male.

Peritrochanteric Pain Syndrome (GTPS)

With a prevalence of 10 to 25%, Greater Trochanteric Pain Syndrome GTPS (women 15% / men 6.6% [\[16\]](#)) is surprisingly one of the most common pain syndromes in orthopedics [\[17\]](#). This may also be related to the fact that it was misinterpreted as trochanteric bursitis for a long time [\[18\]](#), [\[19\]](#) because no connection to a muscular disease was seen. If deep back pain is also present, the prevalence is even 20 to 35% [\[20\]](#).

If one were to also add coxarthrosis, which, according to radiological criteria, affects 14% of people over the age of 55, but only 5% actually show clinical symptoms [\[21\]](#), the number of GTPS cases could be significantly higher. Because at the age of >65 years there are hardly any people who can avoid degenerative changes in the hips [\[22\]](#) and the prevalence varies between 50 and 80% [\[23\]](#), [\[24\]](#), [\[25\]](#).

physiology

The importance of myofascial tissue, which is defined as a complex of muscles (myo) and the surrounding connective tissue (fascia), is of great relevance in orthopedics [\[26\]](#). It is estimated that more than half of all orthopedic patients are affected by myofascial pain syndrome [\[27\]](#), [\[28\]](#). Similar to chronic back pain, which is considered non-specific in 85 to 90% of cases and which is caused by malfunction and tension in the muscles ("myofascial syndrome") [\[29\]](#), [\[30\]](#), [\[31\]](#), [\[32\]](#), expresses itself Inflammation of the pubic bone suspiciously with comparable symptoms.

The hypothesis of an endplate dysfunction is discussed for the myofascial syndrome [\[33\]](#), [\[34\]](#), [\[35\]](#), [\[36\]](#), [\[37\]](#), [\[38\]](#), [\[39\]](#), [\[40\]](#), with the controversial discussions continuing to this day [\[41\]](#), [\[42\]](#), [\[43\]](#). Overloading and traumatic overstretching of a muscle, which cause a subliminal permanent polarization of muscle fibers, are said to be the cause here [\[44\]](#). This causes calcium depletion in the sarcoplasmic reticulum, resulting in local muscle contracture [\[45\]](#), [\[46\]](#), which can be felt as a contracture node (trigger point). The associated local circulatory disturbance ("vascular compression") creates a vicious circle after a spastic muscle stimulates pain receptors, causing it to reflexively contract even more [\[47\]](#). A further complication is the increased energy requirement of the myofascial nodes, which cannot be satisfied under the ischemic conditions [\[48\]](#).

Transferred Pain

In these causal cascades, the "transferred pain" is of particular importance because it explains why local therapeutic efforts often lead to frustrating results. It is postulated [\[49\]](#) that the source of pain is often not to be found in the pain area, but in more or less distant muscles and joints. "Silent synapses" in the posterior horn of the spinal cord ("pain afferents") are said to be activated by the diffusion of neurotransmitters from neighboring neurons, so that the pain unfolds in their projection area [\[50\]](#), [\[51\]](#), [\[52\]](#).

Myofascial Chains

While the term fascia used to be much narrower, today it includes any collagenous tissue. This is interspersed with many mechano- and pain receptors and mediates so-called "anatomy trains": Fascial tissue then provides a direct connection to the individual components of the movement system, so that longitudinal tension and force transmission takes place [\[53\]](#), [\[54\]](#). There are said to be 11 myofascial chains alone, each of which runs on the front and back of the body and sometimes even spirals around the entire body [\[55\]](#). In addition, there is also the concept of a "tensegrity": According to this, the activation of one component always has an impact on neighboring structures, whereby this can be imagined as a flexible connection (myofascium) with solid parts (bones) [\[56\]](#). Muscles therefore do not act separately from one another, but in combination with the help of the facial tissue, so that local changes always result in a mechanical transmission to neighboring body regions.

pubic inflammation

The somewhat misleading addition "inflammation" is based on a bacterial disease of the pubic bone after gynecological or urological operations that is hardly relevant today and describes a painful, non-infectious inflammation of the pubic symphysis or the pubic bone and adjacent structures [\[57\]](#). It is said to be caused by highly

repetitive strain on the adductors of the thigh and possibly the abdominal muscles (m. rectus abdominis). Whether this is due to bone edema (bone bruise) that can only be diagnosed by means of MRI is rather doubtful, since in such cases it is usually a direct injury (e.g. cruciate ligament, ankle) and such an injury is not necessarily associated with pain [\[58\]](#).

If it is the case - as one study claims - that myofascial trigger points, e.g. of the gluteus maximus and medius, primarily produce local pain in the gluteal and sacral region, those of the gluteus minimus muscle on the lower extremities or ankles and those of the pelvic floor muscles mainly in the Pelvic region [\[59\]](#), it seems likely that both hardening at the muscle insertions and pain that is transmitted through myofascial chains play a role here. In this respect, osteitis pubis does not differ from insertion tendinopathy and can therefore also be aptly described as “tennis elbow” of the pubic bone [\[60\]](#). In any case, the symptoms consist of groin pain and pain during isometric adduction of the leg with the hip joint extended or flexed (lift-up test).

Pelvic Pain

Chronic pelvic pain (“longer than 6 months”) or CPPS includes symptoms such as dyspareunia (coital pain), vaginismus, coccygodynia, pain due to pelvic ring instability, anorectal pain and possibly also vulvodynia. Since 78% of interstitial (nonbacterial) cystitis also have a myofascial-related CPP [\[61\]](#), [\[62\]](#), [\[63\]](#), a suspected defect in the glycosome layer (GAG) of the urothelium does not always appear to be the real cause. There is obviously also a connection to endometriosis [\[64\]](#), especially since affected women often also have deep dyspareunia [\[65\]](#), [\[66\]](#), [\[67\]](#). Dyspareunia, which describes pain during penetration, is basically divided into superficial (vaginal introitus) and deep (vaginal canal or pelvic region) [\[68\]](#). Endometriosis, in turn, is defined as endometrial tissue outside the uterus and affects 10% of women of reproductive age [\[69\]](#). So it actually seems almost “logical” if there is also pelvic pain in addition to other symptoms such as dysmenorrhea and painful defecation (dyschezia) [\[70\]](#), especially since the deep dyspareunia is also expressed in interstitial cystitis or bladder pain [\[71\]](#).

Everything seems to be connected to everything. This fuels the assumption that myofascial contractures are the concomitant or perhaps a basis for many LUTS symptoms. The guidelines of the EAU reflect the assessment that female CPPS is a complex neuromuscular-psychosocial process [\[72\]](#), [\[73\]](#), [\[74\]](#). Half of all women who seek help for their CPPS have a history of sexual, physical, or emotional trauma. And a third can be diagnosed with post-traumatic stress disorder (PTSD). ask about the medical history [\[75\]](#).

If you want to go even deeper, myofascial pain occurs more frequently in people who are subject to certain psychosocial factors such as anxiety or stress [\[76\]](#), [\[77\]](#). Just as anxiety and stress can aggravate back pain [\[78\]](#), [\[79\]](#), chronic polyarthritis, soft tissue rheumatism or fibromyalgia [\[80\]](#), [\[81\]](#), chronic pelvic pain also seems to be closely related to a psychological genesis stand [\[82\]](#). In the end, the “knotting” (trigger points) of the muscles could also be viewed as a material-physical enactment of unsolved problems.

Peritrochanteric pain syndrome

Because of the lack of signs of inflammation, the pain syndrome previously treated as bursitis (trochanteric bursitis) was renamed GTPS (Greater Trochanteric Pain Syndrome) [83], [84], [85], [86]. In most cases, a tendinopathy from the gluteal area (M. gluteus medius or minimus) is behind it, although the actual cause remains unclear [87]. The pain usually runs laterally or dorsally to the greater trochanter in the direction of the gluteal muscles and sometimes extends to the lateral thigh and even the knee. There is pronounced pressure pain and an increase in symptoms after exertion.

QRS Pelvicenter rPMS effect

A number of studies have already been indexed on the effectiveness of rPMS in myofascial pain syndromes [88] ("trapezius muscle [89], [90], [91], vastus lateralis muscle after TEP [92], back pain [93], neck muscles [94], other local muscle pain [95], [96] ") although the underlying mechanism of action still remains unclear [97]. Statements such as "pain relief, myostimulation, myorelaxation, decongestant effect or an improvement in blood flow" [98] say little about why a muscle contracture with nodule formation should regress under rPMS use.

The assessment that central mechanisms ("proprioceptive afferents") play a role in pain regulation [99], [100] also ignores the pathogenesis of myofascial trigger points. Possibly no distinction is made here between muscle pain and a malfunction of the peripheral or central nervous system [101]. Studies on rPMS in chronic back pain [102], [103], [104], for example, speak of immediate pain relief in 3 out of 4 studies.

Acute back pain should also respond to rPMS with immediate pain relief, which is also lasting [105]. However, this cannot be explained solely by a gate control theory discussed here. It may be assumed that a muscle contracture follows the impulse of an additive strengthening contraction, ie that relaxation can only be initiated through active tension. This approach is based on experiments on isometric muscle contraction against resistance that produced immediate pain relief in 94% of subjects and sustained pain relief in 63% [106]. This is also confirmed by another study, according to which, for example, a myofascial trigger point can be better treated with electrical stimulation than with a local anesthetic [107].

Scope of treatment and duration of therapy

Derived from the study situation for pelvic pain syndrome in men, 8 to 10 treatments are to be regarded as sufficient. There is disagreement about the frequency, since the gate control theory also plays a role here. However, it is important to remember that muscle tetanus (frequency > 20 Hz) serves to strengthen the muscle [108], [109], while a frequency < 20 Hz is the better choice to induce a single muscle twitch or muscle relaxation [110]. This results in the recommendation to use a frequency of 15 Hz to 20 Hz.

expectation of success

In terms of pain relief (CPPS), rPMS has consistently produced significant results. Unfortunately, the stimulus configurations used in the studies ignore the pathophysiology of a myofascial syndrome. With the correct frequency setting and sequence, therapy optimization can be expected, which would contribute to a significantly better result.

study situation

Only studies on pelvic pain syndrome in men are indexed for rPMS in pelvic pain syndromes. This is not surprising, at least for women with pelvic pain syndrome, since there are some more complex causes behind it. Nevertheless, the men's studies can be used for reference.

Study 1: prospective, randomized and placebo-controlled double-blind study [\[111\]](#)

21 men with an average age of 47.8 years (25 to 67 years) received an rPMS application twice a week (15 minutes at 10 Hz / 15 minutes at 50 Hz) over a period of 4 weeks (8 treatments).

Result:

The mean symptom score had decreased significantly in the active group at both 3 months and 1 year ($p < 0.05$). The best result was achieved for pain symptoms. There was no change in the placebo group.

Study 2: 46 men with CPPS with or without urination problems [\[112\]](#)

46 men with CPPS with or without urinary problems who had not previously responded to any drug therapy were treated with a total of 12 rPMS applications within 6 weeks.

Result:

At 6 months, NIH CPSI score decreased from baseline 25.0 ± 6.9 to 15.6 ± 7.7 , pain score decreased from 11.8 ± 3.7 to 6.9 ± 4.7 (all $p < 0.05$). With regard to benefit, satisfaction and motivation, $> 70\%$ gave a positive response in all subdomains in the BSW Questionnaire (Benefit, Satisfaction and Willingness). Patient satisfaction PPSI (Patient Perception of Symptoms Improvement) was also good after 6 months.

Study 3: 51 patients with LUTS and CPPS [\[113\]](#)

51 patients with LUTS and CPPS received a total of 8 rPMS applications (40 Hz for 10 minutes / 2 minutes rest / 50 Hz for 10 minutes).

Result:

The therapy was completed by 40 patients without absenteeism. 25 of the 40 patients (62.5%) achieved a 30% improvement in their LUTS symptoms. The NIH-CPSI score (NIH-Chronic Prostatitis Symptom Index) decreased from 22.4 to 15.6 ($p < 0.05$). The pain score fell from 9.0 to 6.2, voiding problems from 6.7 to 4.8, quality of life from 6.6 to 4.5. In addition, 11 of 26 patients (42.3%) had at least a 15% improvement in sexual performance. Here the HEF score (International Index of Erectile Function) improved from 44.1 ± 13.5 to 51.6 ± 11.3 ($p < 0.05$).

Study 4: prospective study with 14 CCPS patients [\[114\]](#)

14 CCPS patients received rPMS applications twice weekly over a period of 4 weeks.

Result:

The mean scores (NIH-CPSI) of 4 patients were evaluated. Both the total score ($p < 0.01$) and the sub-area pain or discomfort ($p < 0.02$) decreased significantly. However, there was no significance for the score for micturition ($p = 0.20$). The quality-of-life score increased slightly ($p=0.05$).

Study 5: 30 CPPS patients (average age 39.3 years) [\[115\]](#)

30 CPPS patients (Ø age 39.3 years) received an average of 10.5 rPMS applications with 10 minutes at 40 Hz, pause 2 minutes, 10 minutes at 50 Hz (on-off 5 s: 5 s).

Result:

Twenty of the 30 patients (66.7%) had at least a 30% improvement in symptoms. The total NIH-CPSI score improved significantly from 23.7 to 15.2 ($p < 0.01$). The pain score fell from 11.37 to 6.77, voiding problems from 5.03 to 3.27 and the QOL score from 7.27 to 5.17. In the patients who responded positively to the treatment, the achieved result was maintained in 72% of the patients even 3 months after the end of the therapy.

Study 6: prospective, randomized double-blind study [\[116\]](#)

20 men (Ø 47 years) with CPPS who had failed drug therapy were treated with rPMS for 4 weeks (twice a week) (15 minutes at 10 Hz, 15 minutes at 50 Hz). To cover up the ineffectiveness of the sham device, the sound of the active device was played.

Result:

At 3 months (follow-up), 62% of patients had responded successfully to treatment - vs 13% in the placebo group. After 1 year, the previously achieved result was still valid for 57% of the active and 20% of the placebo group. The pain score (VAS) was 22.7/50 in the active group and fell by 50% to 11.4/50 at 1 year. There was no change in the placebo group (20.4/50 baseline) or increased to 24/50 after one year.

summary

It is becoming increasingly clear that behind most pain syndromes of the musculoskeletal system (back, arthroses, tendinopathies) and thus also of the pelvis there is a myofascial development. Inflammation, cartilage degradation in a joint or dehydration of the intervertebral disc are therefore to be seen as a consequence and not as the cause of the associated restriction of movement or poor posture. In a differentiated consideration of a myofascial syndrome of the pelvis, the cause of the "pubic bone inflammation", which only occurs in athletes, may be a protective reflex

reaction after overloading the adductors or an abdominal muscle inserting into the pubic bone.

Previous childbirth may also play a role in women's pelvic pain syndrome. However, if one also ignores psychosocial reasons, a CPPS or a peritrochanteric pain syndrome - similar to a tennis elbow - can also arise out of "nothing". If the dominant pathology of a myofascial syndrome consists of muscle hardening or contraction blockage due to interspersed painful trigger points, this region is the therapeutic target area. It can be assumed that so-called myofascial function chains exist, which reach from the muscle insertions of the occipital bones (Os occipitale) or from the jaw joint down to the ankle, making the location of the pain origin questionable. On the other hand, there is evidence that the majority of pelvic pain syndromes are closely related to the muscles and fascia of the pelvic floor [\[117\]](#).

Unfortunately, to date there is only one study on rPMS in pelvic pain syndromes on CPPS in men. Due to the universality of myofascial syndromes, however, it seems legitimate to transfer them to women and other myofascial-related indications of the pelvis. However, the usual frequency settings for stress and urge incontinence are not to be used here, but a stimulus configuration that is suitable for inducing muscular relaxation via individual twitches.

Sources

- [1] Dahan R. Rehabilitation of muscle-tendon injuries to the hip, pelvis and groin areas. *Sports Med Arthrosc Rev.* 1997;5:326-333
- [2] Westlin N. Groin pain in athletes from southern Sweden. *Sports Med Arthrosc Rev.* 1997;5:280-284
- [3] Johnson R. Osteitis pubis. *Curr Sports Med Rep* 2003; 2(2):98-102
- [4] Batt ME, McShane JM, Dillingham MF. Osteitis pubis in collegiate football players. *Med Sci Sports Exerc.* 1995; 27(5): 629-633
- [5] Rodriguez C, Miguel A, Lima H et al. Osteitis pubis syndrome in the professional soccer athlete: a case report. *J Athl Train* 2001; 36(4): 437-440
- [6] McMurtry CT, Avioli LV. Osteitis pubis in an athlete. *Calcif Tissue Int.* 1986;38(2):76-77
- [7] Lovell G. The diagnosis of chronic groin pain in athletes: a review of 189 cases. *Aust J Sci Med Sport.* 1995;27(3):76-79
- [8] Pham DV, Scott KG. Presentation of osteitis and osteomyelitis pubis as acute abdominal pain. *Perm J* 2007; 11(2): 65-68
- [9] Holt MA, Keene JS, Graf BK et al. Treatment of osteitis pubis in athletes. Results of corticosteroid injections. *Am J Sports Med* 1995; 23(5): 601-606
- [10] Radic R, Annear P: Use of pubic symphysis curettage for treatment-resistant osteitis pubis in athletes. *Am J Sports Med.* 2008; 36(1):122-128
- [11] Weber O, Kabir K, Müller M et al. Sterile osteitis pubis in a tennis player: diagnosis and treatment options. *German journal sports medicine.* 2010; 61(7-8): 171-174
- [12] Kavrouidakis E, Karampinas PK, Evangelopoulos DS et al. Treatment of osteitis pubis in non-athlete female patients. *Open Orthop J.* 2011; 5:331-334
- [13] Jansen JA, Mens JM, Backx FJ et al. Diagnostics in athletes with long-standing groin pain. *Scand J Med Sci Sports.* 2008; 18(6): 679-690
- [14] Johnson R: Osteitis pubis. *Curr Sports Med Rep.* 2003; 2(2):98-102
- [15] Ahangari A. Prevalence of chronic pelvic pain among women: an updated review. *Pain Physician.* 2014; 17(2): E141-E147
- [16] Segal NA, Felson DT, Torner JC et al. Greater trochanteric pain syndrome: epidemiology and associated factors. *Arch Phys Med Rehab.* 2007; 88:988-992
- [17] Segal NA, Felson DT, Torner JC et al. Greater trochanteric pain syndrome: epidemiology and associated factors. *Arch Phys Med Rehab.* 2007; 88:988-992
- [18] Strauss EJ, Nho SJ, Kelly BT Greater trochanteric pain syndrome. *Sports Med Arthrosc Rev.* 2010;18:113-119
- [19] Barnhouse NC, Wente TM, Voos JE Greater trochanteric pain syndrome: endoscopic treatment options. *Oper Tech Sports Med.* 2012;20:320-324
- [20] Tortolani PJ, Carbone JJ, Quartararo LG. Greater trochanteric pain syndrome in patients referred to orthopedic spine specialists. *Spine J.* 2002; 2: 251-254
- [21] Hackenbroch MH. coxarthrosis. *Orthopäde* 1998;27, 659-667
- [22] Niehard F, Pfeil J. *Orthopädie*. 4th completely revised edition, Thieme Verlag-.2004. p. 484
- [23] Lausmann CS. Localization of pain in advanced coxarthrosis. Dissertation. TU Munich, 2013
- [24] Bagge E, Bjelle A, Eden S et al. Osteoarthritis in the elderly: clinical and radiological findings in 79 and 85 year olds. *Ann Rheum Dis.* 1991; 50, 535-539
- [25] Willaeschus W, Wald A, Swoboda B et al. *rheumatology.* 1996; 21, 2-9
- [26] Giamberardino MA, Affaitati G, Fabrizio A et al. (2011). Myofascial pain syndromes and their evaluation. *Best Pract Res*

Clin Rheumatol 2011;25(2): 185–198

- [27] Wilke J. The importance of myofascial functional chains for the musculoskeletal system with special consideration of mechanical force transfer. dissertation. JWG University Frankfurt. 2016
- [28] Fleckenstein J, Zaps D, Rüger LJ et al. Discrepancy between prevalence and perceived effectiveness of treatment methods in myofascial pain syndrome: results of a cross-sectional, nationwide survey. BMC Musculoskelet Disord. 2010;11:32
- [29] Leinmüller R. Back pain - Most of it is myofascial. Dtsch Ärzteztbl 2008; 105(31-32): A-1657 / B-1430 / C-1397
- [30] Deyo RA. Diagnostic evaluation of LBP: reaching a specific diagnosis is often impossible. Arch Intern Med. 2002;162:1444–1447
- [31] Barmer GEK Healthcare Update 2012. Contributions and analyzes
- [32] Emergency Department patient information. Royal Berkshire NHS Foundation
- [33] Wheeler AH. Myofascial pain disorders: theory to therapy. Drugs 2004; 64(1): 45-62
- [34] Rivner MH. The neurophysiology of myofascial pain syndrome. Curr Pain Headache Rep. 2001 Oct;5(5):432-40
- [35] Partanen JV, Ojala TA, Arokoski JPA. Myofascial syndrome and pain: a neurophysical approach. pathophysiology. 2010; 17(1): 19-28
- [36] Chowdhury N, Oms VI, Goldstein L. Diagnosis and management of myofascial pain syndrome. PPM 2012; 12(2)
- [37] Simons DG, Travell JG. Myofascial Pain and Dysfunction. The trigger point manual. second edition. Lippincott Williams & Wilkins, Baltimore
- [38] Mense S, muscle tone and muscle pain. ManMed 2005; 43(3): 156-161
- [39] Hong CZ, Somons DG. Pathophysiologic and electrophysiologic mechanisms of myofascial trigger points. Arch Phys Med Rehab 1998; 79: 863-872
- [40] Hah JP, An in vivo microanalytical technique for measuring the local biochemical milieu of human skeletal muscle. Appl Physiol 2005; 99: 1977-1988
- [41] Quintner JL, Bove GM, Cohen ML. A critical evaluation of the trigger point phenomenon. Rheumatology (>Oxford): 2015; 54(3): 392-399
- [42] Dommerholt J, Gerwin RD. A critical evaluation of Quintner et al: missing the point. J Bodyw Mov Ther. 2015; 19(2): 193-204
- [43] Quintner JL, Bove GM, Cohen ML. Response to Dommerholt and Gerwin: Did we miss the point ? J Bodyw Mov Ther 2015, 19(3): 394-395
- [44] Beat de Jung. Muscular joint pain. Diagnostic and therapeutic aspects. Ars Medici 2010; 11:444-447
- [45] Simons DG, Travell JG. Myofascial origins of low back pain. 1. Principles of diagnosis and treatment. Postgrad Med 1983 Feb; 73(2): 66, 68 -70
- [46] Beat de Jung. Muscular joint pain. Diagnostic and therapeutic aspects. Ars Medici 2010; 11: 444-447
- [47] Opitz G. Muscle pain. Part 1: Theory of muscle pain and acupuncture therapy. pain acupuncture. 2005; 3:151-163
- [48] Simon's DG. Myofascial pain syndromes: where are we ? Where are we going? Arch Phys Med Rehab 1988; 69(3 Pt 1): 207-12
- [49] Mense S. Muscle tone & muscle pain. Manual medicine. 2005; 43(3):156-161
- [50] Friction JR. Myofascial pain: clinical characteristics and diagnostic criteria. J Musculoskeletal Pain. 1993; 12:37-47
- [51] Mense S. The basis for muscular spasm. Abstracts of NZAMSM. Annual Scientific Conference. Queenstown, NZ 2003
- [52] Mense S. Biochemical pathogenesis of myofascial pain. J Musculo-skeletal Pain. 1996; 4:145-162
- [53] Myers TW. The "anatomy trains". Part 1. J Bodyw Mov Ther 1997; 1:91-101
- [54] Myers TW. The "anatomy trains". Part 2. J Bodyw Mov Ther 1997; 1: 135-145
- [55] Krause F, Wilke J. Relevance of myofascial chains in movement therapy of musculoskeletal diseases. Critical literature overview. Manual Therapy 2017; 21: 189-194
- [56] Ingber DE. The architecture of life. Sci Am. 1998; 278(1): 48-57
- [57] Hopp S Osteitis pubis. German journal for sports medicine. 2008; 59: 100-101
- [58] Krüger J. Bone edema in the os pubis - suggested staging using MRI. Sports ortho trauma, 2012; 28:182-188
- [59] Simons DG, Travell JG. Simons DG, Travell JG. Myofascial origins of low back pain. 3. Pelvic and lower extremity muscles. Postgrad Med 1983 Feb; 73(2): 99-108
- [60] Klingelhöffer W. Is footballer's pubic osteitis homemade? Sports kinesiological methods as a supplement to conventional medicine. COMED, August 2013
- [61] Bassaly R, Tidwell N, Bertolino S et al. Myofascial pain and pelvic floor dysfunction in patients with interstitial cystitis. International Urogynecology Journal. 2010;22(4):413–418
- [62] Doggweiler-Wiygul R, Wiygul JP. Interstitial cystitis, pelvic pain, and the relationship to myofascial pain and dysfunction: A report on four patients. WorldJ Urol. 2002;20(5):310–314
- [63] Itza F, Zarza D, Serra L et al. Myofascial pain syndrome in the pelvic floor: a common urological condition. Actas Urologicas Espanolas. 2010;34(4):318–326
- [64] Orr NL, Noga H, Williams C. Deep dyspareunia in endometriosis: role of the bladder and pelvic floor. J Sex Med 2018; 15(8): 58-1166
- [65] Shum LK, Bedaiwy MA, Allaire C et al. Deep dyspareunia and sexual quality of life in women with endometriosis. Sex Med 2018; 6:224-233
- [66] Lukic A, Di Properzio M, De Carlo S et al. Quality of sex life in endometriosis patients with deep dyspareunia before and after laparoscopic treatment. Arch Gynecol Obstet 2016; 293:583-590
- [67] Fritzer N, Tammaa A, Haas D et al. When sex is not on fire: a prospective multicentre study evaluating the short-term effects of radical resection of endometriosis on quality of sex life and dyspareunia. Eur J Obstet Gynecol Reprod Biol 2016; 197:36-40
- [68] Yong PJ, Sadownik L, Brotto LA. Concurrent deepsuperficial dyspareunia: prevalence, associations, and outcomes in a multidisciplinary vulvodynia program. J Sex Med 2015;12:219-227
- [69] Eskenazi B, Warner ML. Epidemiology of endometriosis. Obstet Gynecol Clin North Am 1997;24:235-258
- [70] Ferrero S, Esposito F, Abbamonte LH et al. Quality of sex life in women with endometriosis and deep dyspareunia. Fertil Steril 2005;83:573-579
- [71] Yong PJ. Deep dyspareunia in endometriosis: a proposed framework based on pain mechanisms and genito-pelvic pain penetration disorder. Sex Med Rev 2017;5:495-507
- [72] Engeler DS, Baranowski AP, Dinis-Oliveira P et al. European Association of Urology. The 2013 EAU guidelines on chronic pelvic pain: is management of chronic pelvic pain a habit, a philosophy, or science ? 10 years of development. EUR euros. 2013; 64(3): 431-439
- [73] Engeler DS,, Baranowski AP, Borvicka J et al. European Association of Urology. Guidelines on chronic pelvic pain
- [74] Potts JM, Payne CK. Urologic chronic pelvic pain. Pain. 2012; 153(4): 755-758

- [75] Meltzer-Brody S, Leerman J, Zolnoun D et al. Trauma and posttraumatic stress disorder in women with chronic pelvic pain. *Obstet Gynecol* 2007; 109(4): 902-908
- [76] Klotz SGR, Ketels G, Loewe B et al. Myofascial findings and psychopathological factors in patients with chronic pelvic pain syndrome. *Pain Med* 2018 May 19. doi. 10.1093/pm/pny097. Epub ahead of print.
- [77] Ohrbach R, Michelotti A. The role of stress in the etiology of oral parafunction and myofascial pain. *Oral Maxillofac Surg Clin North Am.* 2018; 30(3): 369-379
- [78] Shaw WS, Hartvigsen J, Woiszwilllo MJ et al. Psychological distress in low back pain: a review of measurement scales and levels of distress reported in the first 2 months after pain onset. *Arch Phys Med Rehab* 2016; 97(9): 1573-1587
- [79] Hoy D, Brooks P, Blyth F et al. The epidemiology of low back pain. *Best Practice Res Clin Rheumatol* 2010; 24(6): 769-781
- [80] Hermann M, Schölmerich J, Straub RH. Stress and rheumatic diseases. *Rheum Dis Clin North Am.* 2000; 26(4): 737-763
- [81] Herrmann M, Schölmerich J, Straub RH. Stress and rheumatic diseases. *Rheum Dis Clin North Am* 2000; 26(4): 737-763
- [82] Haugsted GK, Kirste U, Leganger S et al. Somatocognitive therapy in the management of chronic gynecological pain. A review of the historical background and results of a current approach. *Scand J Pain.* 2018; 2(3): 124-129
- [83] Williams BS, Cohen SP. Greater trochanteric pain syndrome: a review of anatomy, diagnosis and treatment. *Anesth Analg.* 2009;108:1662-1670
- [84] Shbeeb MI, O'Duffy JD, Michet CJ et al. Evaluation of glucocorticosteroid injection for the treatment of trochanteric bursitis. *J Rheumatol.* 1996;23:2104-2106
- [85] Strauss EJ, Nho SJ, Kelly BT Greater trochanteric pain syndrome. *Sports Med Arthrosc Rev.* 2010;18:113-119
- [86] Barnhouse NC, Wente TM, Voos JE. Greater trochanteric pain syndrome: endoscopic treatment options. *Oper Tech Sports Med.* 2012;20:320-324
- [87] Genth B, von Düring M, von Engelhardt LV et al. Analysis of the sensory innervations of the greater trochanter for improving the treatment of greater trochanteric pain syndrome. *Clin Anat* 2012;25:1080-1086
- [88] Beaulieu LD, Schneider C. Repetitive peripheral magnetic stimulation to reduce pain or improve sensorimotor impairments: a literature review on parameters of application and afferents recruitment. *Neurophysiol Clin.* 2015; 45(3): 223-234
- [89] Sollmann N, Albers L, Jung NH et al. Magnetic stimulation of the upper trapezius muscles in patients with migraine - a pilot study. *Science Direct.* 2016; 20(6):888-897
- [90] Smania N, Corato E, Fiaschi A et al. Repetitive magnetic stimulation. A novel approach for myofascial pain syndrome. *J Neurol* 2005; 252:307-314
- [91] Smania N, Corato E, Fiaschi A et al. Therapeutic effect of peripheral repetitive magnetic stimulation on myofascial pain syndrome. *Clin Neurophysiol.* 2003; 114(2):350-358
- [92] Baek J, Park N, Lee B et al. Effects of repetitive peripheral magnetic stimulation over vastus lateralis in patients after hip replacement surgery. *Ann Rehabilitation Med* 2018; 42(1): 67-75
- [93] Lim YH, Song JM, Choi EH et al. Effects of repetitive peripheral magnetic stimulation on patients with acute low back pain: a pilot study. *Ann Rehabilitation Med* 2018; 42(2): 229-238
- [94] Smania N, Corato E, Fiaschi A et al. Repetitive magnetic stimulation. A novel approach for myofascial pain syndrome. *J Neurol* 2005; 252:307-314
- [95] Pujol J, Pascual-Leone A, Dolz C et al. The effect of repetitive magnetic stimulation on localized musculo-skeletal pain. *neuro report.* 1998; 9(8): 1745-1748
- [96] Beaulieu LD, Schneider C. Effects of repetitive peripheral magnetic stimulation on normal or impaired motor control. A review. *Neurophysiol Clin.* 2013; 43(4): 251-260
- [97] Smania N, Corato E, Fiaschi A et al. Repetitive magnetic stimulation. A novel approach for myofascial pain syndrome. *J Neurol* 2005; 252: 307-314
- [98] Zarkovic D, Kazalakova K. Repetitive peripheral magnetic stimulation as pain management solution in musculoskeletal and neurological disorders - a pilot study. *Int J Physiother.* 2016; 3(6): 671-675
- [99] Melzack R, Wall PD. Pain mechanisms: a new theory. *Science* 1965; 150: 971-979
- [100] Stanton-Hicks M, Salamon J. Stimulation of the central and peripheral nervous system for the control of pain. *J Clin Neurophysiol.* 1997; 14: 46-62
- [101] Baron R. Peripheral neuropathic pain: from mechanisms to symptoms. *Clin J Pain.* 2000; 16(2): S12-S20
- [102] Lo YL, Fook-Chong S, Huerto AP, George JM. A randomized, placebo-controlled trial of repetitive spinal magnetic stimulation in lumbosacral spondylotic pain. *Pain Med* 2011;12:1041-1045
- [103] Masse-Alarie H, Flamand VH, Moffet H, Schneider C. Peripheral neurostimulation and specific motor training of deep abdominal muscles improve posturo-motor control in chronic low back pain. *Clin J Pain.* 2013;29:814-823
- [104] Masse-Alarie H, Beaulieu LD, Preuss R, Schneider C. Repetitive peripheral magnetic neurostimulation of multifidus muscles combined with motor training influences spine motor control and chronic low back pain. *Clin Neurophysiol.* 2017;128:442-453
- [105] Lim YH, Song JM, Choi EH et al. Effects of repetitive peripheral magnetic stimulation on patients with acute low back pain: a pilot study. *Ann Rehabilitation Med* 2018; 42(2): 229-238
- [106] Lewit K, Simons DG. Myofascial pain: relief by post-isometric relaxation. *Arch Phys Med Rehab.* 1984; 65(8): 452-456
- [107] Ga H, Koh HJ, Choi JH et al. Intramuscular and nerve root stimulation vs lidocaine: injection to trigger points in myofascial pain syndrome. *J Rehabilitation Med* 2007; 39(5): 463-467
- [108] Han TR, Shin HI, Kim IS. Magnetic stimulation of the quadriceps femoris muscle: comparison of pain with electrical stimulation. *At J Phys Med Rehab* 2006; 85(7): 593-599
- [109] Ito T, Tsubahara A, Watanabe S. Use of electrical or magnetic stimulation for generating hip flexion torque. *Am J Phys Med Rehabil* 2013;92(9):755-761
- [110] Beaulieu LD, Schneider C. Effects of repetitive peripheral magnetic stimulation on normal or impaired motor control. A review. *Neurophysiol Clin.* 2013; 43(4): 251-260
- [111] Rowe E, Smith C, Laverick L et al. A prospective, randomized, placebo-controlled, double-blind study of pelvic electromagnetic therapy for the treatment of chronic pelvic pain syndrome with 1 year follow-up. *J Urol* 2005; 173(6): 2044-2047
- [112] Kim TH, Han DH, Cho WJ et al. The efficacy of extracorporeal magnetic stimulation for treatment of chronic prostatitis / chronic pelvic pain syndrome patients who do not respond to pharmacotherapy. *Urology.* 2013; 82(4):894-898
- [113] Park HR, Gho SG, Kim HJ. Therapeutic efficacy of extracorporeal magnetic therapy on low urinary tract symptoms and sexual function in Chronic Pelvic Pain Syndrome. *Korean J Urol.* 2006; 47(6): 645-650
- [114] Kim SW, Kim SH, Lee CH et al. Clinical efficacy of extracorporeal magnetic innervation for chronic pelvic pain syndrome. *Korean J Androl.* 2003; 21:44-47
- [115] Lee KC, Choi H, Park HS et al. Therapeutic efficacy of extracorporeal magnetic therapy in chronic pelvic pain syndrome.

Korean J Urol. 2003; 44: 693-696

[116] Patel A, Rowe E, Leverick L. A prospective, randomized, placebo-controlled, double-blinded study of electromagnetic therapy (ExMI) in the treatment of chronic pelvic pain syndrome in men. Presented at the XVth Congress of the European Association of Urology Geneva, Switzerland, April 7-10, 2001

[117] Simons DG, Travell JG. Myofascial origins of low back pain. 3. Pelvic and lower extremity muscles. Postgrad Med 1983; 73(2): 99-105, 108
